

Prevalence of Seasonal Variation of Haemorrhagic stroke more in winter than summer and its common risk factors.

Ahmed S¹, Islam MA², Bari A³, Hasan MJ⁴, Pathan AS⁵, Rubel MM⁶, Sarker CN⁷

Abstract

The present descriptive cross-sectional study was conducted in the department of medicine, Community Based Medical College Hospital Bangladesh over a period of one year during March 2013 to February 2014. The study was conducted to describe Haemorrhagic stroke during summer and winter. An attempt was also made to observe in the frequency of common risk factors of stroke by seasons. A total 146 patients of any age irrespective of sex fulfilling the WHO criteria of acute stroke and confirmed by CT Scan were selected from consecutive admission in the Department of Medicine, Community Based Medical College Hospital Bangladesh. Detail history and through clinical examinations were done. Routine and relevant investigations were carried out. The mean age of the patients was 60.9±14.3 years. A male preponderance was observed in the study. In summer 65.8% of patients and in winter 34.2% patients were presented. Ischemic stroke was present in 52.1% patients and 47.9% patients had haemorrhagic stroke. The study found that the frequency of ischaemic stroke during summer (65.8%) was significantly greater than that during winter (34.2%). The frequency of haemorrhagic stroke during winter (65.8%) was significantly greater than that during summer (39.6%). Hypertension was the most important risk factor and other risk factors were smoking, Diabetes mellitus, tobacco chewing, ischaemic heart disease, dyslipidaemia, oral contraceptive pill, alcohol consumption, atrial fibrillation and past history of stroke. Increasing age was also noted as a risk factor (60.7% > 60 years). Most of the risk factors were homogeneously distributed between two seasons and between ischaemic and haemorrhagic group. Hypertension was significantly higher in haemorrhagic stroke patients compared to ischaemic stroke patients.

CBMJ 2016 January; Vol. 05 No. 01 P: 9-13

Key words: Ischaemic stroke, Haemorrhagic stroke Summer, Winter.

Introduction

After coronary artery disease and cancer stroke is the third commonest cause of death in the developed countries¹. It is a common medical emergency. The incidence rises steeply with age, and in many lower and middle income countries. It is rising in association with less healthy lifestyles. About one fifth of patients with an acute stroke die within a month of the event and at least half of those who survive are left with physical disability². In the under developed world, with better control of infectious diseases and with increase in life expectancy. Thus stroke remains as a major cause of mortality, morbidity and great burden to the family members, society and hospital manpower. Non modifiable risk factors of stroke include age, sex, family history, race and ethnicity. Modifiable risk factors are hypertension, diabetes mellitus, cardiac disease (particularly

atrial fibrillation), Hyper lipidaemia, smoking, transient ischemic attacks, asymptomatic carotid artery stenosis, alcohol abuse and physical inactivity³. Prevalence of stroke may

1. *Dr. Sultan Ahmed
Assistant Professor, Department of Medicine,
Community Based Medical College Bangladesh.
2. Dr. Md. Aminul Islam
Associate Professor, Department of Medicine,
Community Based Medical College Bangladesh.
3. Dr. Mohammad Abdul Bari
Assistant Professor, Department of Medicine,
Community Based Medical College Bangladesh.
4. Dr. Mahmud Javed Hasan
Assistant Professor, Department of Nephrology,
Community Based Medical College Bangladesh.
5. Dr. Md. Ashraf Siddique Pathan
Assistant Professor, Department of Psychiatry
Community Based Medical College Bangladesh.
6. Professor Dr. Chanda Nendu Bhushan Sarker
Professor & Head, Department of Medicine,
Community Based Medical College Bangladesh.
7. Dr. Mostufa Mahmud Rubel
Assistant Registrar, Department of Medicine,
Community Based Medical College Bangladesh.

*Address of correspondence

Email : drsultan1975@gmail.com
Mobile : 01717 316654

be related to its risk factors. So, the study was planned to prospectively haemorrhagic stroke more in winter with common risk factors.

Method

This descriptive cross sectional study was conducted in Community Based Medical College Hospital, Bangladesh over a period of one year from March 2013 to February 2014. Patients admitted in medicine wards with acute (<7 days) development of focal or global disturbance of neurological function with or without previous history of stroke and confirmed by CT scan of brain were included in the study. Patients who, could not avail CT scan of brain or who died within 24 hours of hospitalization were excluded from the study. Consecutive purposive sampling technique was followed for selection of cases and total 146 patients were included from the consecutive admission in medicine wards.

Data were collected using semi-structured questionnaire containing all the variables of interest. Data were recorded through face to face interview with nearest relative/attendant physical examination findings recorded by investigator after a thorough clinical examination, CT Scan report collected on the same day or when done. Collected data were compiled, processes and analyzed with the help of computer using SPSS (statistical package for social science version 11.5). Both descriptive and inferential statistics were used. Descriptive statistics were used for grouping and organizing of demographic variable and inferential (chi-square and fisher exact test) statistics were used to see the association between seasons and types of stroke.

The word Health Organizations definition of stroke as the rapidly developing clinical symptoms and or signs of focal (at times global) disturbance of cerebral function with symptoms lasting more than 24 hours or leading to death with no apparent cause other than that of vascular origin was applied for selection of cases. According to Bangladesh Meteorological Department, Dhaka, the

season of Bangladesh is broadly divided into summer and winter depending on variation of temperature. The duration of summer is from March to November and that of winter is from December to February⁴. This definition was applied in our study. However, to remain the frequency difference due to duration we convert frequency into a per month rate.

Data were analyzed using chi-square test.

Result

A total 146 stroke patients admitted during March 2013 to February 2014 were included and CT Scan was done to confirm the diagnosis of stroke and its types. The finding of the study obtained from data analysis are shown on table below.

Table-I: Distribution of study subjects by demographic variable (N=146)

Variables	Number	%
Age (Years)		
<40	10	6.8%
40-50	22	15%
50-60	25	17%
60-70	40	27.4%
≥ 70	49	33.6%
Sex		
Male	85	58.2%
Female	61	41.8%

I Mean age (60.9 ± 14.3) years, Range=(20-90)

Table-II: Distribution of subjects by season of occurred and types of stroke (N=146)

Seasons	Total stroke	Ischaemic stroke	Haemorrhage
Summer	96 (65.8%)	58 (60.4%)	38 (39.6%)
Winter	50 (34.2%)	18 (36%)	32 (64%)
Total	146 (100%)	76 (52.1%)	70 (47.9%)

Table- III: Monthly frequency of stroke during summer and winter

Types of stroke			
Season	Ischaemic (N=76)	Haemorrhagic (N=70)	P-Value
Summer	11 (14.5%)	8 (11.5%)	0.732
March- 2013	5 (6.6%)	2 (2.9%)	0.098
April- 2013	5 (6.6%)	3 (4.3%)	0.389
May- 2013	4 (3.3%)	3 (4.3%)	0.772
June- 2013	5 (6.6%)	3 (4.3%)	0.389
July- 2013	5 (6.6%)	3 (4.3%)	0.389
August- 2013	11 (14.5%)	7 (10%)	0.640
September- 2013	4 (5.3%)	3 (4.3%)	0.772
October- 2013	8 (10.6%)	6 (8.6%)	0.772
November-2013			
Winter			
December 2013	10 (13.2%)	18 (23.7%)	0.040
January- 2014	4 (5.3%)	7 (10%)	0.030
February-2014	4 (5.3%)	7 (10%)	0.030

Table IV: Frequency of ischemic and haemorrhagic stroke during summer and winter.

Season			
Season	Total stroke	Ischaemic stroke	Haemorrhage
Ischaemic	58	18 (36%)	
Haemorrhagic	38	32 (64%)	< 0.001

Chi- square (X^2) test was employed to analyze the data (chi-square (X^2) Value = 15.9, df=1)

Table V: Distribution of subjects by risk factors (N=146)

Risk factors	Number	%
Hypertension	100	68.5%
Smoking	46	31.5%
Alcohol	02	1.4%
Tobacco chewing	15	10.3%
Oral contraceptive	12	8.2%
Diabetes	24	16.4%
IHD	04	2.7%
AF	05	3.4%
Past H/O Smoking	02	1.4%

Table VI: Comparison of risk factors between ischaemic and haemorrhagic stroke.

Group			
Risk factors	Ischemic (N=76)	Haemorrhagic (N=70)	P Value
Hypertension	40 (52.6%)	60 (85.7%)	< 0.001
Smoking	31 (40.8%)	15 (21.4%)	0.163
Alcohol	1 (1.3%)	1 (1.4%)	0.625
Tobacco chewing	8 (10.5%)	7 (10%)	0.757
Oral contraceptive	6 (7.9%)	6 (8.6%)	0.687
Diabetes	16 (21.1%)	8 (11.4%)	0.091
IHD	3 (3.9%)	1 (1.4%)	0.562
AF	4 (3.3%)	1 (1.4%)	0.150
Past H/O Stroke	1 (1.3%)	1 (1.4%)	0.375

Data were analyzed using chi-square test, fisher exact test.

Table VII: Comparison of risk factors between summer and winter

Season			
Risk factors	Summer N= 96	Winter (N=50)	P Value
Hypertension	65 (67.5%)	31 (62%)	0.580
Smoking	31 (32.3%)	12 (24%)	0.221
Alcohol	01 (01%)	01 (2%)	0.483
Tobacco chewing	06 (6.3%)	14 (28%)	< 0.001
Oral contraceptive	07 (7.3%)	04 (8%)	0.772
Diabetes	16 (6.7%)	04 (8%)	0.044
IHD	00	03 (6%)	0.014
AF	01 (01%)	02 (4%)	0.83
Past H/O Smoking	03 (3%)	01 (2%)	0.715

Data was analyzed using chi-square test. Fisher exact test was done to analyze the data.

Table VIII: Distribution of subject by lipid profile (N=146)

Lipid Profile	Number	Percentage
Total cholesterol (mg/dl)		
Normal <200	113	77.4%
Raised ≥ 200	33	22.6%
HDL Cholesterol		
Normal ≥ 40 mg/dl	86	58.9%
Low < 40mg/dl	59	40.4%
LDL Cholesterol (mg/dl)		
Normal <100	74	50.7%
Raised ≥ 100	72	49.3%
Triglycerides (mg/dl)		
Normal < 150	89	60.9%
Raised ≥ 150	56	38.4%

References

1. Mackay J, Menash G. The atlas of heart disease and stroke, world health organization, Geneva 2000.
2. Langhorre P. Stroke disease walker BR, College NR, Rueston SH. Penman 1D. Davidsons principle and practice of medicine, churchill living stone. New Delhi. An imprint of Else vier 2014; P1231-47.
3. Sacco RL, Benjamin EJ, Broderick JP, Dyken M, Easten JD et al. Risk factors. Stroke 1997; 28:1507-17.
4. Hannan MA. Study of seasonal variation of stroke. MD thesis. University of Dhaka, Dhaka, Bangladesh 1999.
5. Sang- Pil Yun, Woo-Sang Jung, Sang-K Wan, Moon, Ki-Ho Cho, Young-Suk, Kim, Hyung-Sup, Bae, Seasonal variation of Acute stroke Hospital Based Study J Korean Oriental Med. 2006; 29 (2):1-6
6. Haque A, Mannan MA. Rahman H2. Prognosis of acute stroke. Bangladesh of Neuro Science 1987; 3: 42-7
7. Anwarullah AKM, Habib M, Mohammad QD, Ahmed S, Nahar S, Review of risk factors for stroke: Study of 100 cases. Bangladesh J of Neuroscience 1993; 9:11
8. Jovanovic Z. Risk factors for stroke In young people Srp Arh Celok Lek 1996; 124, 21.15
9. Masood K, Maqsood MA, Qudri AA. Stroke: Seasonal variation In Etiology, Professional med. 2006; 13(4): 680-6
10. Rohwell PM, Wroe SJ, Statterly J, Warlow CP. Is stroke incidence related to season or temperature? Lancet, 1996; 347: 934-936.
11. Goldstein LB, Adams R, Alberts MJ, Appel LJ, Brass LM, Bushnell CD, Primary Prevention of Ischemic stroke; A guideline from the am Heart Association/American stroke Association stroke council cosponsored by the atherosclerotic peripheral vascular disease interdisciplinary warving group; Cardiovascular nursing council; Clinical cardiology council; Nutrition, Physical activity and metabolism council; and the quality of care and outcomes research interdisciplinary warving group. The american Academy of Neurology affirms the value of this guideline stroke 2006; 37:1583-1633
12. Liu LH, Chia LG. The effects of hypertension, Diabetes mellitus, atrial fibrillation. Transient ischemic attack and smoking on stroke in Chinese people. Chung Hua. Hsueh TS chin Teupei, 1991; 47:1105
13. Thrift AG, McNeil JJ, Fxbes A, Donnan GA, Risk factors for cerebral haemorrhage in the era of well controlled hypertension; Melbourne risk factor study (MERFS) group stroke. 1996, 27:2020-2025.
14. Miah AH. Study of seasonal variation in types of stroke and its common risk factors resident physical (medium) MMCH, Dhaka, Bangladesh, 2012.